

The abundance of HOC^+ In diffuse clouds

H. Liszt¹ and R. Lucas² and J. H. Black³

¹ National Radio Astronomy Observatory, 520 Edgemont Road, Charlottesville, VA, USA 22903-2475

² Institut de Radioastronomie Millimétrique, 300 Rue de la Piscine, F-38406 Saint Martin d'Hères, France

³ Chalmers Centre for Astrophysics and Space Science, Onsala Space Observatory, Chalmers University of Technology, 43992 Onsala, Sweden

received February 2, 2008

Abstract. We used the Plateau de Bure Interferometer to search for $\lambda 3\text{mm}$ absorption lines of HOC^+ from local diffuse and translucent clouds occulting compact extragalactic mm-wave continuum sources. We detected HOC^+ in three directions with column densities only 70-120 times below that of the HCO^+ isomer, a factor 5-50 higher than typically found in dense dark gas but comparable to recent observations of dense photon-dominated regions. The observed amounts of HOC^+ $N(\text{HOC}^+)/N(\text{H}_2) = 3 - 6 \times 10^{-11}$ can be made in **quiescent** diffuse gas at thermal gas-kinetic rates if the $\text{H}_2\text{O}/\text{OH}$ ratio is of order unity, in mild violation of extant observational limits.

Key words. interstellar medium – molecules

1. Introduction.

The roster of polyatomic molecules known to exist in diffuse interstellar gas has been steadily enlarged over the last fifteen years and now numbers nearly one dozen. Two of these, C_3 (Maier et al., 2001; Roueff et al., 2002; Oka et al., 2003) and H_3^+ (McCall et al., 2003) were recently discovered in optical and near-infrared absorption spectra. The others have accumulated gradually over the last twenty or so years in cm-wave and mm-wave radiofrequency absorption data and now include H_2CO (Federman & Willson, 1982; Nash, 1990; Marscher et al., 1993; Liszt & Lucas, 1995; Moore & Marscher, 1995), HCO^+ (Lucas & Liszt, 1996), NH_3 (Nash, 1990), C_2H (Lucas & Liszt, 2000), 3h2 (Cox et al., 1988; Lucas & Liszt, 2000), HCN and HNC (Liszt & Lucas, 2001) and H_2S and HCS^+ (Lucas & Liszt, 2002).

Observationally, the polyatomics observed with radio techniques are often seen to be strongly and directly related to simpler species observed at optical wavelengths, *i.e.* HCO^+ to OH (Liszt & Lucas, 1996; Lucas & Liszt, 1996) HCN and HNC to CN (Liszt & Lucas, 2001) *etc.* These connections point to fundamental *lacunae* in our understanding of the chemistry of even the simplest species in diffuse clouds: **simple** models which reproduce the abundances of diatomics, however easily, invariably fall orders of magnitude short of explaining the related polyatomics. Conversely, the high abundances of some polyatomics transform other mysteries; the poorly-understood abundances of CO and CS in diffuse clouds are readily understandable given the recombination of the observed columns of

HCO^+ (Liszt & Lucas, 2000) and HCS^+ (Lucas & Liszt, 2002) respectively.

The work described here rounds out to an even dozen the number of polyatomics known to exist in diffuse and translucent clouds. HOC^+ , the energetically-disfavored HCO^+ isomer whose 89.4874 GHz $J=1-0$ transition we sought, was indeed detected along several of the sightlines studied in the other molecular species mentioned in the preceding paragraphs. The HCO^+ - HOC^+ comparison is especially interesting now because both species have recently been detected over a wide range of physical conditions in dense gas: our new spectra reinforce an emerging view which relates molecular abundances in dense and diffuse PDR. Section 2 gives details of the observations, which are discussed in Sect. 3 and summarized in Sect. 4.

2. Observations of HOC^+

The 89.4874 GHz $J=1-0$ transition of HOC^+ was observed at the Plateau de Bure at various times during the period 2000-2001 toward the five sources listed in Table 1. Profiles were taken with 39.1 kHz (0.13 km s^{-1}) channel separation and 70 kHz resolution and subsequently once hanning smoothed, yielding spectra with the line/continuum rms values as listed in Table 1.

The column density in the lowest ($J=0$) level of a simple linear molecule is related to the integrated optical depth of the $J=1-0$ transition as

$$N_0 = \frac{8.0 \times 10^{12} \text{ cm}^{-2} \int \tau_{10} dv}{\mu^2 (1 - \exp(-h\nu_{10}/kT_{\text{exc}}))} \quad (1)$$

Table 1. Background sources observed

Source	l	b	$\sigma_{l/c}^1$	$\int \tau(\text{HOC}^+) dv$ km s ⁻¹
B0355+508	150.38°	-1.60°	0.0022	0.042(0.005)
B0415+379	161.68°	-8.82°	0.0029	0.051(0.004)
B0528+134	191.37°	-11.01°	0.0101	<0.036
B1730-130	12.03°	+10.81°	0.0081	<0.026
B2200+420	92.13°	-10.40°	0.0041	0.017(0.006)

¹ rms noise in line/continuum ratio at 140 kHz (0.46 km s⁻¹) resolution**Table 2.** Column densities

Source	N(HOC ⁺) 10 ¹⁰ cm ⁻²	N(HCO ⁺) 10 ¹² cm ⁻²	N(OH) 10 ¹³ cm ⁻²
B0355+508	9.01(1.07)	6.51(0.02)	14.87(0.67)
B0415+379	10.9(0.90)	12.8(0.40) ²	42.50(0.10)
B0528+134	< 7.7 ¹	2.28(0.02)	4.72(0.31)
B1730-130	< 5.7 ¹	1.25(0.02)	1.34(0.08)
B2200+420	3.7(1.3)	2.58(0.03)	5.73(0.21)

¹ limits are 2 σ ² Entry is 59 $\pm$ 2 N(H¹³CO⁺)**Table 3.** Column density ratios¹

Source	N(HCO ⁺)/N(HOC ⁺)	N(HOC ⁺)/N(OH)
B0355+508	72(9)	$6.1(0.7) \times 10^{-4}$
B0415+379	117(12)	$2.6(0.3) \times 10^{-4}$
B0528+134	>30	< 1.6×10^{-3}
B1730-130	> 18	< 4.3×10^{-3}
B2200+420	70(24)	$6.5(2.1) \times 10^{-4}$

¹ limits are 2 σ


where μ is the permanent dipole moment of the molecule, T_{exc} is the excitation temperature of the J=1-0 transition and ν_{10} is its frequency.

With the exception of CO (Liszt & Lucas, 1998), the molecules observed at mm-wavelengths in diffuse gas are well-described by assuming that the rotational ladder is thermalized at the temperature of the cosmic microwave background, in which case the total column density can be calculated straightforwardly from observations of one transition. We take the dipole moments of HOC⁺ and HCO⁺ as 2.8D and 3.93D, respectively, a slight departure from prior work in which we used 4.07D for HCO⁺. Given these assumptions, one has $N(X)/\int \tau_{10} dv = 2.146 \times 10^{12} \text{ cm}^{-2}$ for X=HOC⁺, $1.093 \times 10^{12} \text{ cm}^{-2}$ for X=HCO⁺ and $1.128 \times 10^{12} \text{ cm}^{-2}$ for X=H¹³CO⁺.

3. Observed and expected abundances of HOC⁺ in diffuse gas

3.1. Observed abundance of HOC⁺

The mm-wave absorption data are summarized in Tables 1-3 and line profiles toward two sources are shown in Figures 1 and 2. The HCO⁺ and OH column densities are taken from our

**Fig. 1.** HOC⁺ and H¹³CO⁺ observed toward B0415+379 (aka 3C111)

prior published (Lucas & Liszt, 1996; Liszt & Lucas, 1996) and unpublished work; for B0415+379 (see Fig. 1) we assert $N(\text{HCO}^+) = 59 \pm 2 N(\text{H}^{13}\text{CO}^+)$ (Lucas & Liszt, 1998) because of the high optical depth in the main isotope. HOC⁺ is found to have relative abundances $\text{HCO}^+:\text{HOC}^+ = 70\text{--}120$, $\text{HOC}^+:\text{OH} = 3 - 6 \times 10^{-4}$ toward three sources (see Table 3). The results obtained toward the other two, B0528 and B1730, are not of great significance. The relative abundance of OH, for which it is found that $X(\text{OH}) = n(\text{OH})/n(\text{H}_2) = 1.0 \pm 0.2 \times 10^{-7}$ along the four sightlines where both $N(\text{H}_2)$ and $N(\text{OH})$ have been measured (Liszt & Lucas, 2002), sets the relative abundance scale for HOC⁺, $X(\text{HOC}^+) = 3 - 6 \times 10^{-11}$.

Note that the OH column density toward B2200+420 is only 20% larger than that seen along the archetypical diffuse line of sight toward ζ Oph, $N(\text{OH}) = 4.7 \pm 0.7 \times 10^{13} \text{ cm}^{-2}$ (Van Dishoeck & Black, 1986; Roueff, 1996), while the reddening ($E_{B-V} = 0.33$ mag toward B2200+420 from Schlegel et al. (1998)) is the same. The larger total OH column density seen toward B0355+508 at low galactic latitude (-1.6°) is roughly evenly divided among 5 kinematically-separated diffuse clouds of modest OH and HCO⁺ column density, only two of which have prominent CO emission (Lucas & Liszt, 1996; Liszt & Lucas, 1996). The line of sight toward B0415+379 is somewhat darker (see the discussion in Lucas & Liszt (1998)) but relatively little of the carbon along even this line of sight is in CO and the density is fairly low, as gauged by the weakness of emission from species beside CO (Lucas & Liszt, 1996).




Fig. 2. HOC^+ and HCO^+ observed toward B2200+420 (aka BL Lac)

3.2. Expected abundance of HOC^+

The most obvious path to HOC^+ in diffuse gas is *via* the reaction $\text{C}^+ + \text{H}_2\text{O} \rightarrow \text{HOC}^+ + \text{H}$, which proceeds with a rate constant $1.8 \times 10^{-9} \text{ cm}^3 \text{ s}^{-1}$ in the UMIST reaction database¹; these two reactants also produce HCO^+ at half the rate, but this is not an important source of HCO^+ . We parametrize the unknown abundance of water relative to OH and assume that all free carbon is once ionized in a fully molecular gas having a free carbon abundance $\text{N}(\text{C})/\text{N}(\text{H}) = 1.4 \times 10^{-4}$ (Savage & Sembach, 1996). In these terms, the volume formation rate of HOC^+ from the reaction of C^+ and water can be expressed as $\text{dn}(\text{HOC}^+)/\text{dt} = 5.04 \times 10^{-20} \text{ n}(\text{H}_2)^2 \text{ X}(\text{H}_2\text{O})/\text{X}(\text{OH}) \text{ cm}^{-3} \text{ s}^{-1}$.

Both HCO^+ and HOC^+ are formed from the reaction of $\text{H}_3^+ + \text{CO} \rightarrow (\text{HCO}^+ \text{ or } \text{HOC}^+) + \text{H}_2$. The $\text{HCO}^+/\text{HOC}^+$ branching ratio is typically taken as 20 (Illies et al., 1982) or more (the current UMIST reaction database has a branching ratio of 62). The volume formation rate from this reaction may be written as $\text{dn}(\text{HOC}^+)/\text{dt} = 2.7 \times 10^{-23} \text{ n}(\text{H}_2)^2 (\text{X}(\text{H}_3^+) \text{ X}(\text{CO})/10^{-12}) \text{ cm}^{-3} \text{ s}^{-1}$. The relative abundance of CO is known along these lines of sight to be 1-5% of the free carbon abundance (Liszt & Lucas, 1998), *i.e.* $1 - 5 \times 10^{-6}$. The observed amounts of H_3^+ , while surprisingly large (McCall et al., 2003) nevertheless still yield $\text{X}(\text{H}_3^+) < 10^{-6}$. Thus the only circumstance in which the reaction of H_3^+ and CO could be competitive in forming HOC^+ would be for very small water abundances, in which case neither formation route would suffice to form the observed HOC^+ .

The reaction of CO^+ and H_2 forms only HCO^+ , contrary to the claim of Fuente et al. (2003) (their Sect. 6.1).

HOC^+ is likely destroyed by two processes which proceed with very nearly equal rates in diffuse gas. The HOC^+ isomer lies some 17,000K higher in energy compared to HCO^+ and interconversion by molecular hydrogen ($\text{HOC}^+ + \text{H}_2 \rightarrow \text{HOC}^+ + \text{H}_2$) has recently been shown to proceed with a rate constant $k_I = 4 \times 10^{-10} \text{ cm}^3 \text{ s}^{-1}$ nearly independent of temperature between 25K and 300 K (Smith et al., 2002). This is 70% smaller than in the UMIST database, enhancing the prospects for reproducing $\text{X}(\text{HOC}^+)$. The destruction rate for isomerization can be written as $\text{dn}(\text{HOC}^+)/\text{dt} = -4 \times 10^{-10} \text{ n}(\text{H}_2) \text{ n}(\text{HOC}^+) \text{ cm}^{-3} \text{ s}^{-1}$.

Dissociative recombination, $\text{HOC}^+ + e^- \rightarrow \text{CO} + \text{H}$, proceeds with a rate constant $1.1 \times 10^{-6} (30/T_K) \text{ cm}^3 \text{ s}^{-1}$. Assuming that free carbon is once-ionized and expressing the electron fraction in terms of the ratio of free carbon to H_2 expected for a fully molecular diffuse gas (*i.e.* $\text{N}(\text{C})/\text{N}(\text{H}_2) = 2 \times 1.4 \times 10^{-4}$ (Savage & Sembach, 1996)). The destruction rate from recombination can be written as $\text{dn}(\text{HOC}^+)/\text{dt} = -3.1 \times 10^{-10} \text{ n}(\text{H}_2) \text{ n}(\text{HOC}^+) (30/T_K) (\text{X}(\text{e})/\text{X}(\text{C}^+)) \text{ cm}^{-3} \text{ s}^{-1}$. The expected electron abundance is a small multiple of the free carbon abundance (Liszt, 2003) so dissociative recombination and interconversion probably compete quite effectively to destroy HOC^+ .

The net destruction rate from both processes is of order $-10^{-9} \text{ n}(\text{H}_2) \text{ cm}^{-3} \text{ s}^{-1}$ and the expected abundance of HOC^+ is $\text{X}(\text{HOC}^+) = 5 \times 10^{-11} \text{ X}(\text{H}_2\text{O})/\text{X}(\text{OH})$ or $\text{X}(\text{HOC}^+)/\text{X}(\text{OH}) = 5 \times 10^{-4} \text{ X}(\text{H}_2\text{O})/\text{X}(\text{OH})$. The observed amounts of HOC^+ can be made in diffuse gas if the abundance of water is comparable to that of OH. Limits on the OH/water ratio in diffuse clouds $\text{N}(\text{OH})/\text{N}(\text{H}_2\text{O}) > 2-3$ were derived by Van Dishoeck & Black (1986) toward ζ Oph and ζ Per.

Given the weakness of the limits on the water abundance in diffuse clouds, straightforward gas-phase chemistry fails only weakly to produce the observed amount of HOC^+ . This situation can be compared to the case of HCO^+ where the thermal reaction of C^+ upon the known quantities of OH falls 30-50 times short of producing the observed values of $\text{N}(\text{HCO}^+)$ via the reaction chain $\text{C}^+ + \text{OH} \rightarrow \text{CO}^+ + \text{H}$, $\text{H}_2 + \text{CO}^+ \rightarrow \text{HCO}^+ + \text{H}$, $\text{HCO}^+ + e^- \rightarrow \text{CO} + \text{H}$. At thermal rates, this series of reactions (which predicts an HCO^+/OH ratio dependent only on temperature) produces $\text{N}(\text{HCO}^+)$ scarcely larger than the observed $\text{N}(\text{HOC}^+)$. The predicted abundance of HCO^+ could be increased in quiescent gas by hypothesizing that it recombines slowly, but only at the cost of understanding CO (Liszt & Lucas, 2000).

Clearly, knowledge of the abundance of water is the limiting factor in assessment of the diffuse cloud chemistry of HOC^+ .

3.3. Comparison with HCO^+ and HOC^+ in dense dark and bright gas

The work presented here is the first evidence of HOC^+ in diffuse gas, but even the case for its existence in the dense molecular ISM was marginal before the pioneering work of Apponi and his collaborators during a brief period in the mid-1990's while the NRAO 12m telescope was still operating.

¹ available at <http://www.rate99.co.uk/>

To summarize the prior results we note that the $\text{HCO}^+/\text{HOC}^+$ ratio in dense dark gas is typically as large as 1000-9000 (Ziurys & Apponi, 1995; Apponi & Ziurys, 1997), but is found to be much smaller (50-400) in dense PDR like the Orion Bar (Apponi et al., 1999) and/or NGC7023 (Fuente et al., 2003). In these PDR the relative abundance of HOC^+ is $X(\text{HOC}^+) = 1 - 3 \times 10^{-11}$, quite similar to the diffuse clouds studied here.

The difference between the $\text{HCO}^+/\text{HOC}^+$ ratios seen in diffuse and dark dense gas is largely the result of the low ionization fraction in the latter. Whereas HOC^+ is destroyed only slightly faster than HCO^+ in diffuse gas, the isomerization reaction undergone by HOC^+ is much faster than recombination when the electron fraction is 10^{-7} instead of 10^{-4} . In dense dark gas HCO^+ and HOC^+ are formed by $\text{H}_3^+ + \text{CO} \rightarrow \text{H}_2 + (\text{HCO}^+ \text{ or } \text{HOC}^+)$ with a branching ratio 20-60:1 in favor of HCO^+ and this ratio is increased by another factor of 100 by the unequal destruction rates. Under favorable conditions, measured $\text{HCO}^+/\text{HOC}^+$ ratios could be good diagnostics of the ionization fraction in dense gas.

Reasoning by analogy from the discussion of diffuse and dark gas, we infer that the relatively low $\text{HCO}^+/\text{HOC}^+$ ratios in dense PDR also reflect the relative weakness of isomerization. In cases where isomerization is weak both isomers are destroyed by recombination at roughly equal rates, and their abundance ratio is more directly indicative of the relative rates at which the two isomers are formed.

4. Summary

We have added HOC^+ to the roster of polyatomic molecules known to exist in diffuse clouds. The $\text{HCO}^+/\text{HOC}^+$ ratio, 70-120, is much smaller than in dense dark clouds but quite comparable to the most extreme values seen in dense PDR. The inferred relative abundance $X(\text{HOC}^+) = 3 - 6 \times 10^{-11}$, similar to that seen in dense PDR, can be synthesized at thermal rates in quiescent diffuse gas if the water/OH ratio is of order unity. This is a mild discrepancy, much less severe than is inferred for HCO^+ , as the water/OH ratio is currently **known only** to be less than 0.3-0.5.

Several authors have recently noted that hydrocarbons like 3h2 and C_2H have unexpectedly high abundances in dense PDR (Fosse et al., 2002; Fuente et al., 2003; Teyssier et al., 2004), very much as has been observed in diffuse clouds (Lucas & Liszt, 2000). The $\text{HCO}^+/\text{HOC}^+$ ratios found here **in diffuse clouds, which are PDR of lower density and presumably weaker illumination (hence similar ionization rates per H_2), constitute another example of chemical similarities between dense and diffuse PDR.** It will be interesting to see how much more complete a parallel can be drawn between the two regimes.

Acknowledgements. The National Radio Astronomy Observatory is operated by AUI, Inc. under a cooperative agreement with the US National Science Foundation. IRAM is operated by CNRS (France), the MPG (Germany) and the IGN (Spain). We owe the staff the Plateau de Bure our thanks for their assistance in taking the data. We thank the referee for helpful comments.

References

- Apponi, A. J., Pesch, T. C., & Ziurys, L. M. 1999, *ApJ*, 519, L89
- Apponi, A. J. & Ziurys, L. M. 1997, *ApJ*, 481, 800
- Cox, P., Guesten, R., & Henkel, C. 1988, *A&A*, 206, 108
- Federman, S. R. & Willson, R. F. 1982, *ApJ*, 260, 124
- Fosse, D., Teyssier, D., Gerin, M., Pety, J., Abergel, A., & Habart, E. 2002, in *Chemistry as a Diagnostic of Star Formation*, 104-107
- Fuente, A., Rodriguez-Franco, A., Garcia-Burillo, S., Martin-Pintado, J., & Black, J. H. 2003, *A&A*, 406, 899
- Illies, A. J., Jarrold, M. F., & Bowers, M. T. 1982, *J. Chem. Phys.*, 77, 5847
- Liszt, H. 2003, *A&A*, 398, 621
- Liszt, H. & Lucas, R. 2001, *A&A*, 370, 576
- , 2002, *A&A*, 391, 693
- Liszt, H. S. & Lucas, R. 1995, *A&A*, 299, 847
- , 1996, *A&A*, 314, 917
- , 1998, *A&A*, 339, 561
- , 2000, *A&A*, 355, 333
- Lucas, R. & Liszt, H. 1998, *A&A*, 337, 246
- Lucas, R. & Liszt, H. S. 1996, *A&A*, 307, 237
- , 2000, *A&A*, 358, 1069
- , 2002, *A&A*, 384, 1054
- Maier, J. P., Lakin, N. M., Walker, G. A. H., & Bohlender, D. A. 2001, *ApJ*, 553, 267
- Marscher, A. P., Moore, E. M., & Bania, T. M. 1993, *ApJ*, 419, L101
- McCall, B. J., Huneycutt, A. J., Saykally, R. J., Geballe, T. R., Djuric, N., Dunn, G. H., Semaniak, J., Novotny, O., Al-Khalili, A., Ehlerding, A., Hellberg, F., Kalhori, S., Neau, A., Thomas, R., Österdahl, F., & Larsson, M. 2003, *Nature*, 422, 500
- Moore, E. M. & Marscher, A. P. 1995, *ApJ*, 452, 671
- Nash, A. G. 1990, *Astrophys. J., Suppl. Ser.*, 72, 303
- Oka, T., Thorburn, J. A., McCall, B. J., Friedman, S. D., Hobbs, L. M., Sonnentrucker, P., Welty, D. E., & York, D. G. 2003, *ApJ*, 582, 823
- Roueff, E. 1996, *Mon. Not. R. Astron. Soc.*, 279, L37
- Roueff, E., Felenbok, P., Black, J. H., & Gry, C. 2002, *A&A*, 384, 629
- Savage, B. D. & Sembach, K. R. 1996, *Ann. Rev. Astrophys. Astron.*, 34, 279
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- Smith, M. A., Schlemmer, S., von Richthofen, J., & Gerlich, D. 2002, *ApJ*, 578, L87
- Teyssier, D., Fossé, D., Gerin, M., Pety, J., Abergel, A., & Roueff, E. 2004, *A&A*, 417, 135
- Van Dishoeck, E. F. & Black, J. H. 1986, *Astrophys. J., Suppl. Ser.*, 62, 109
- Ziurys, L. M. & Apponi, A. J. 1995, *ApJ*, 455, L73